

1. Product Rule: $y = (x^2 + 1)(x^3 + 1) \Rightarrow$

$$y' = (x^2 + 1)(3x^2) + (x^3 + 1)(2x) = 3x^4 + 3x^2 + 2x^4 + 2x = 5x^4 + 3x^2 + 2x.$$

Multiplying first: $y = (x^2 + 1)(x^3 + 1) = x^5 + x^3 + x^2 + 1 \Rightarrow y' = 5x^4 + 3x^2 + 2x$ (equivalent).

2. Quotient Rule: $F(x) = \frac{x - 3x\sqrt{x}}{\sqrt{x}} = \frac{x - 3x^{3/2}}{x^{1/2}} \Rightarrow$

$$F'(x) = \frac{x^{1/2} \left(1 - \frac{9}{2}x^{1/2}\right) - (x - 3x^{3/2}) \left(\frac{1}{2}x^{-1/2}\right)}{(x^{1/2})^2} = \frac{x^{1/2} - \frac{9}{2}x - \frac{1}{2}x^{1/2} + \frac{3}{2}x}{x} = \frac{\frac{1}{2}x^{1/2} - 3x}{x} = \frac{1}{2}x^{-1/2} - 3$$

Simplifying first: $F(x) = \frac{x - 3x\sqrt{x}}{\sqrt{x}} = \sqrt{x} - 3x = x^{1/2} - 3x \Rightarrow F'(x) = \frac{1}{2}x^{-1/2} - 3$ (equivalent).

For this problem, simplifying first seems to be the better method.

13. $y = (r^2 - 2r)e^r \xrightarrow{\text{PR}} y' = (r^2 - 2r)(e^r) + e^r(2r - 2) = e^r(r^2 - 2r + 2r - 2) = e^r(r^2 - 2)$

14. $y = \frac{1}{s + ke^s} \xrightarrow{\text{QR}} y' = \frac{(s + ke^s)(0) - (1)(1 + ke^s)}{(s + ke^s)^2} = -\frac{1 + ke^s}{(s + ke^s)^2}$

21. $y = 2xe^x \Rightarrow y' = 2(x \cdot e^x + e^x \cdot 1) = 2e^x(x + 1)$. At $(0, 0)$, $y' = 2e^0(0 + 1) = 2 \cdot 1 \cdot 1 = 2$, and an equation of the tangent line is $y - 0 = 2(x - 0)$, or $y = 2x$. The slope of the normal line is $-\frac{1}{2}$, so an equation of the normal line is $y - 0 = -\frac{1}{2}(x - 0)$, or $y = -\frac{1}{2}x$.


23. (a) $y = f(x) = \frac{1}{1 + x^2} \Rightarrow$

$$f'(x) = \frac{(1 + x^2)(0) - 1(2x)}{(1 + x^2)^2} = \frac{-2x}{(1 + x^2)^2}.$$

So the slope of the

tangent line at the point $(-1, \frac{1}{2})$ is $f'(-1) = \frac{2}{2^2} = \frac{1}{2}$ and its

equation is $y - \frac{1}{2} = \frac{1}{2}(x + 1)$ or $y = \frac{1}{2}x + 1$.



29. $f(x) = \frac{x^2}{1 + x} \Rightarrow f'(x) = \frac{(1 + x)(2x) - x^2(1)}{(1 + x)^2} = \frac{2x + 2x^2 - x^2}{(1 + x)^2} = \frac{x^2 + 2x}{x^2 + 2x + 1} \Rightarrow$

$$f''(x) = \frac{(x^2 + 2x + 1)(2x + 2) - (x^2 + 2x)(2x + 2)}{(x^2 + 2x + 1)^2} = \frac{(2x + 2)(x^2 + 2x + 1 - x^2 - 2x)}{[(x + 1)^2]^2}$$

$$= \frac{2(x + 1)(1)}{(x + 1)^4} = \frac{2}{(x + 1)^3},$$

so $f''(1) = \frac{2}{(1 + 1)^3} = \frac{2}{8} = \frac{1}{4}$.